

# gravity waves from quantum stress tensor fluctuations in inflation

Jen-Tsung Hsiang  
National Dong-Hwa University, Taiwan

# introduction

- inflation beautifully explains what the universe looks like today,
- but it is not clear what actually drives inflation,
- nor about how long it lasts, as long as inflation is over 60 e-foldings,
- it implies that the scales we are interested right now may derive from transplanckian scales in the beginning of inflation,
- this observation may offer a tool to probe transplanckian physics.

# introduction

- we will look into some of these issues by examining backreactions of the matter on the gravity waves.
- other than (active) quantum fluctuations of the metrics, there is an additional (passive) component induced by quantum fluctuations of matter,
- quantum fluctuations of the matter results in fluctuations of its energy stress tensor,
- in turn, by the Einstein equation, the stress tensor fluctuations drive (passive) metric fluctuations.

# introduction

- it turns out that this induced component of the metric fluctuations may distort the tensor mode power spectrum of the CMB in the high frequency end,
- the extent of correction depends on duration of inflation,
- besides, it contains contributions from the (trans)plackain modes of the matter field,
- so it can used as a testground of ultra-high energy physics, when combined with the future observation data from the PLANCK or LISA-type experiments.

# configuration

consider a spatially flat de Sitter universe


$$ds^2 = a^2(\eta) [-d\eta^2 + d\mathbf{x}^2] .$$

with  $\eta$  the conformal time and  $a(\eta) = -(H\eta)^{-1}$ ,  $\eta < 0$ .

- let  $g_{\mu\nu} = \gamma_{\mu\nu} + h_{\mu\nu}$ , where  $\gamma_{\mu\nu}$  is the background de Sitter metric, and  $h_{\mu\nu}$  is the metric perturbation – tensor modes.
- choose the transverse-tracefree (TT) gauge:

$$h^{\mu\nu}{}_{;\nu} = 0, \quad h = h^\mu{}_\mu = 0, \quad h_{\mu\nu} u^\nu = 0,$$

- “;” denotes covariant derivative wrpt the bkgd metric,
- $u^\nu$  is some timelike vector.



Only a portion of the de Sitter spacetime describes inflationary expansion.

# (active) gravity waves

Lifshitz (1946) showed the tensor modes in a spatially flat universe behave as massless scalars,

$$\square_s h^\mu{}_\nu = 0.$$

The  $\square_s$  is a scalar wave operator.

- gravitons are equivalent to a pair of minimally coupled massless scalar fields.
- the corresponding quantum fluctuations are so-called “active”.
- the power spectrum of the active gravity wave produced during inflation is flat.

# generation of gravity waves

on the other hand, gravity waves may be generated by a source,

$$\square_s h^\mu{}_\nu = -16\pi G_N S^\mu{}_\nu.$$

here  $S^\mu{}_\nu$  is the transverse-tracefree part of the source stress tensor.

in the semiclassical theory, they may be generated by

- the renormalized expectation value of its stress tensor (PRD **83**, 084027).
- quantum fluctuations of the matter (PRD **84**, 103515).



# quantum fluctuations of the stress tensor

integrating linearized Einstein equation

$$\square_s h^\mu{}_\nu = -16\pi G_N S^\mu{}_\nu,$$

by the retarded Green's function

$$\square_s G_R(x, x') = -\frac{\delta(x - x')}{\sqrt{-\gamma}},$$

gives the induced metric perturbation/fluctuation

$$h^\mu{}_\nu(x) = 16\pi G_N \int d^4x' \sqrt{-\gamma'} G_R(x, x') S^\mu{}_\nu(x').$$

we may form the metric correlation function  $K^\mu{}_\nu{}^\rho{}_\sigma$

$$\begin{aligned} & K^\mu{}_\nu{}^\rho{}_\sigma(x, x') \\ &= (16\pi)^2 \int d^4y \sqrt{-\gamma} \int d^4y' \sqrt{-\gamma'} G_R(x, y) G_R(x', y') C^\mu{}_\nu{}^\rho{}_\sigma(y, y'), \end{aligned}$$

in terms of a stress tensor correlation function  $C^\mu{}_\nu{}^\rho{}_\sigma$ , where

$$\begin{aligned} K^\mu{}_\nu{}^\rho{}_\sigma(x, x') &= \langle h^\mu{}_\nu(x) h^\rho{}_\sigma(x') \rangle - \langle h^\mu{}_\nu(x) \rangle \langle h^\rho{}_\sigma(x') \rangle, \\ C^\mu{}_\nu{}^\rho{}_\sigma(x, x') &= \langle S^\mu{}_\nu(x) S^\rho{}_\sigma(x') \rangle - \langle S^\mu{}_\nu(x) \rangle \langle S^\rho{}_\sigma(x') \rangle. \end{aligned}$$

- independent of the renormalization scheme.
- anomaly free for the conformal matter.

- the evolution of metric nonlocally depends on matter (history dependent).
- for a conformally invariant field

$$C_{\mu\nu\rho\sigma}^{FRW}(x, x') = a^{-4}(\eta)a^{-4}(\eta') C_{\mu\nu\rho\sigma}^{Mink}(x, x').$$

- define the power spectrum by the Fourier transform of the equal-time correlation function,

$$P(k) = \int \frac{d^3\mathbf{R}}{(2\pi)^3} e^{i\mathbf{k}\cdot\mathbf{R}} K(\eta = \eta', \mathbf{R})$$

- it is related to the power spectrum in cosmology by  $\mathcal{P}(k) = 4\pi k^3 P(k)$ .

# sudden switching

1st scenario: if quantum fluctuations of matter couple w/ gravity at the onset of inflation  $\eta = \eta_0$ , and then integrate forward in time to the end of inflation at  $\eta = \eta_r$ , then we have

$$\mathcal{P}_s(k) = -\frac{4H^2}{3\pi} S^2 k^2 (1 + k^2 H^{-2}), \quad k|\eta_0| \gg 1,$$

$S$  is the expansion factor during inflation.

- negative power spectra,
- blue tilt  $\mathcal{P}(k) \propto k^4$ ,
- grows as  $S^2$ .

# exponential switching

2nd scenario: the coupling to the fluctuating stress tensor is switched on gradually with a switching function  $e^{\lambda\eta}$ .

note that  $\lambda^{-1}$  is the approximate conformal time at which the interaction begins.

$$\mathcal{P}_\epsilon(k) = -\frac{3H^3}{8\pi} S k (1 + k^2 H^{-2})$$

$S$  is the expansion factor during inflation.

- negative power spectra,
- blue tilt  $\mathcal{P}(k) \propto k^3$ ,
- grows as  $S^1$ .

as for the negative power spectra: the Wiener-Khinchine theorem requires a non-negative spectrum for a regular correlation function.

however, for quadratic quantum operators, such as a stress tensor, the positive definite quantity in this theorem may not exist b/c the corresponding correlation function is highly singular.

this allows for negative power spectra. (Phys. Lett. A**375**, 2296.)

another example: for the flat-space EM energy density

$$P(k) = -\frac{k^5}{960\pi^5}.$$

# numerical estimation - I

- consider perturbations of the order of the present horizon size,  $\ell \approx 10^{61} \ell_p$ .
- WMAP constrain these perturbations to satisfy  $h \leq 10^{-5}$ .
- $|K_{now}| \leq 10^{-10}$ .
- this limits, for exponential switching,

$$S_e < 10^{40} \left( \frac{10^{16} \text{GeV}}{E_R} \right)^7$$

- it is compatible with adequate inflation  $S \geq 10^{23}$  for the flatness problem.
- $\because \mathcal{P} < 0$ , quantum stress tensor fluctuations during inflation tends to produce ANTI-correlated gravity wave fluctuations.

# numerical estimation - II

- consider perturbations of the order of 100 km.
- LIGO constrains these perturbations to satisfy  $h \leq 10^{-24}$ .
- $|K_{now}| \leq 10^{-48}$ .
- we have a bound on  $S$ , for exponential switching,

$$S_e < 10^{25} \left( \frac{10^{13} \text{GeV}}{E_R} \right)^7$$

- compatible with adequate inflation if  $E_R \leq 10^{13} \text{GeV}$ .
- $\because \mathcal{P} < 0$ , stress tensor fluctuations during inflation tends to produce anti-correlated gravity wave fluctuations.



# summary

- the gravity waves are generated by quantum stress tensor fluctuations during inflation.
- this induced gravity waves tend to anti-correlated.
- its power spectra are negative, nonscale-invariant.
- this spectrum also depends on the duration of inflation.
- the effect is in principle observable in that gravity wave modes are no longer exactly solutions of the Lifshitz equation.
- this possibility does require the contribution of modes which were transplanckian at the beginning of inflation.
- if we apply similar considerations to different inflation models/alternative gravity theories, together with observation data from LISA or BBO, it may improve our understanding of inflation/transplanckian physics.